PA007748.h1Fs-b: Biotinylated Recombinant Anti-human CD28 Monoclonal Antibody (Clone: 15E8), Human IgG1 Fc Silent Kappa

Catalog No.: PA007748.h1Fs-b | **Product Name:** Biotinylated Recombinant Anti-human CD28 Monoclonal Antibody, Human IgG1 Fc Silent Kappa (Clone: 15E8)

Biotinylated recombinant anti-human CD28 monoclonal antibody (Clone 15E8), engineered with a Human IgG1 Fc-silent domain to eliminate non-specific Fc receptor binding during flow cytometric analysis. Features site-directed biotinylation via a C-terminal Avi tag on the heavy chain, with variable regions derived from anti-human CD28 clone (15E8) and expressed in mammalian cells.

PRODUCT SPECIFICATIONS

Catalog No.	PA007748.h1Fs-b
Product Name	Biotinylated Recombinant Anti-human CD28 Monoclonal Antibody (Clone: 15E8), Human IgG1 Fc Silent Kappa
Supplier Name	Syd Labs, Inc.
Brand Name	Syd Labs
Clone	15E8
Isotype	Human IgG1 Fc Silent Kappa.
Source/Host	The site-directed biotinylated anti-human CD28 monoclonal antibody (clone: 15E8) was produced in mammalian cells.
Specificity/Sensitivity	The site-directed biotinylated recombinant 15E8 antibody specifically binds to human CD28.
Applications	Flow cytometry (FC) with various dye conjugation through streptavidin; conjugation to streptavidin agarose beads or streptavidin magnetic beads.
Form Of Antibody	0.2 uM filtered solution, pH 7.4, no stabilizers or preservatives.
Conjugate	Biotin-conjugated.
Purity	>95% by SDS-PAGE under reducing conditions.
Shipping & Storage	The 15E8 antibody is shipped with ice pack. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 1 month from date of receipt, 2 to 8°C as supplied.

BACKGROUND & APPLICATION INFORMATION

Recombinant human IgG1 monoclonal antibody with site-directed biotinylation through the Avi tag at the C-terminus of the antibody heavy chain.

The 15E8 antibody binds to human CD28 expressed on T cells and thymocytes. The site-directed biotinylation enables sensitive detection via streptavidin-dye conjugates in flow cytometry, cell surface staining, and T cell co-stimulation studies while avoiding Fc receptor-mediated background signals.